



*Brief report*

## Detection of children's malingering on Raven's Standard Progressive Matrices

R. Kim McKinzey<sup>1\*</sup>, Jörg Prieler<sup>2</sup> and John Raven<sup>3</sup>

<sup>1</sup> Private Practice, Albany, USA

<sup>2</sup> Dr G. Schuhfried GmbH, Hyrtlstraße, Austria

<sup>3</sup> Consultant, Edinburgh, Scotland, UK

A formula for detecting faked Raven's SPM profiles was cross-validated on 44 children and adolescents (ages 7–17). It yielded a false negative rate of 64%. However, a rule using three very easy items (i.e., any of A3, A4 or B1 missed) yielded a hit rate of 95%, with 5% false positive and negative rates. All but two of the participants were able to produce lower scores when asked to fake the test.

Faust (Faust, 1996; Faust, Ziskin, & Hiers, 1991) and Rogers (Rogers, Harrell, & Liff, 1993) have documented the need for detection of neuropsychological malingering. To solve the problem, Gudjonsson and Shackleton (1986) validated a formula using Raven's Standard Progressive Matrices (SPM; J. C. Raven, 1958). The formula has the distinct advantage of being usable on protocols given in the past, as it requires no special administration procedures. The formula was cross-validated (McKinzey, Podd, Krehbiel, & Raven, 1999) on 46 experimental malingerers and 381 people from the adult standardization sample (J. Raven, J. C. Raven, & Court, 2000). The formula yielded a cross-validated 26% false negative rate and a 5% false positive rate.

However, the formula was validated and cross-validated on adults, and one does not have to be an adult to fake a psychological test. Faust has also documented the ability of normal adolescents (Faust, Hart, Guilmette, & Arkes, 1988) and children (Faust, Hart, & Guilmette, 1988) to produce abnormal neuropsychological test results that are believable and undetectable by the same methods that are useful in detecting cortically based neuropsychological deficits.

The formula was therefore applied to a sample of children and adolescents.

\* Requests for reprints should be addressed to R. K. McKinzey, 400 29th St, Ste. 315, Oakland, CA 94609, USA (e-mail: editor@wpe.net).

## Method

### Participants

All 44 participants came from schools in Vienna, Austria, and were all Caucasian. Girls comprised 57% of the sample. Age range was 7-17 years ( $M = 12.5$ ,  $SD = 2.6$ ), with nine in the 7-10 age range, 27 in the 11-14 range, and eight in the 15-17 range. Participants' total score range was 18-59 ( $M = 43.95$ ). Using the US norms (J. Raven, 2000), their percentile range was 24-99 ( $M = 65.9$ ).

### Procedure

Parents signed consent forms. All participants were given Raven's SPM. First, the test was administered according to standard instructions (J. Raven *et al.*, 2000), and then the same participants were asked to take the test again, this time with the instructions (in German):

We know that some people don't try their best on this test. We'd like to find a way to catch them. To help us, please do as badly on this test as you can, without getting caught.

The Raven answers were applied to the formula  $(2A + B) - (D + 2E)$ , where A, B, D and E refer to the number of correct responses in each of the Raven subsets (Gudjonsson & Shackleton, 1986; McKinzey *et al.*, 1999). The result of the formula is termed the 'rate of decay', and is compared to the rate of decay by total score cut-offs (see McKinzey *et al.*, 1999, for details).

## Results

In the faking condition, the participants produced total scores (range 2-59,  $M = 9.6$ ) and percentiles (range 1-99,  $M = 6.3$ ) substantially below that of the normal condition, with a mean difference between the two conditions of 34 ( $t = -14.003$ ,  $p < .0001$ ) and mean intrasubject total score differences of 34 ( $SD = 15.36$ , range 1-57). Not surprisingly, using the formula cut-offs validated for adults proved highly inaccurate, with a false positive rate of 7% and a false negative rate of 64%.

The inaccuracy rates were minimally artificially elevated by the participants' inability to fake the test. Other studies (e.g., Heaton, Smith, Lehman, & Vogt, 1978; McKinzey, Podd, Krehbiel, Mensch, & Trombka, 1997) have found that some participants are unable to fake a given test sufficiently to produce abnormal results, a problem referred to as a 'threat to external validity' (Rogers & Cruise, 1998). Such ineffectual faking attempts are of little consequence in interpretation, since the difference between the true and actual scores will be minimal. However, one of the participants had no difference between the two conditions, and another did one item better! When these two participants were eliminated, the false negative rate dropped by only two percentage points. Changing the cut-off points was the obvious next step, but visual inspection revealed a far more obvious (and accurate) method of detecting the participants in the faking condition.

According to the standard instructions, the testee must agree to the correct answers to the first two items. The third item is very easy, and only one of the participants (an 8-year-old girl who produced an average Raven IQ score) answered it incorrectly. When

these same participants malingered on the test, item A3 was overwhelmingly answered incorrectly. Table 1 presents the classification results of the simple rule that a testee getting item 3 incorrect is showing insufficient effort. The hit rate of the rule is 90%, with a false positive rate of 2% and false negative rate of 17%. Combining the item 3 rule and the rate of decay formula *decreased* the hit rate by 1%.

**Table 1.** Item 3 rule results

	Formula result		Totals
	Raven faked N (% of row)	Raven not faked N (% of row)	
Malingering condition	35 (83)	7 (17)	42 (100)
Normal condition	1 (2)	41 (98)	42 (100)
	36	48	84

Notes: Percentages are rounded. The chi-square statistic is highly significant ( $\chi^2 = 58.37$ ,  $p < .0001$ ; Fisher's Exact  $p < .0001$ ). The table does not include the two participants who did not produce malingered scores.

**Table 2.** Item A3 or A4 or B1 rule results

	Formula result		Totals
	Raven faked N (% of row)	Raven not faked N (% of row)	
Malingering condition	40 (95)	2 (5)	42 (100)
Normal condition	2 (5)	40 (95)	42 (100)
	42	42	84

Notes: Percentages are rounded. The chi-square statistic is highly significant ( $\chi^2 = 68.76$ ,  $p < .0001$ ; Fisher's Exact  $p < .0001$ ). The table does not include the two participants who did not produce malingered scores.

Using analyses of item difficulty (J. Raven *et al.*, 2000) as a guide, items A4 and B1 were similarly identified as increasing the hit rate. Table 2 presents the classification results, removing the two unsuccessful faking participants. Applying the rule that missing any of items A3, A4 or B1 (all extremely easy items) should be interpreted as showing insufficient effort produces a 95% hit rate, with equal false positive and negative rates of 5%. The two false positives were the 8-year-old girl (normal condition total score = 21, 44th percentile) and an 11-year-old girl (normal condition total score = 35, 34th percentile). The two false negatives were a 12-year-old girl (faking condition total score = 19, down from 34) and a 14-year-old girl (faking condition total score = 34, down from 53).

## Discussion

There are few methods of detecting faked IQ test results. Other cross-validated methods are available to detect malingering of neuropsychological deficits—for example, the Test of Malingered Memory (TOMM), a commercial product designed to detect

neuropsychological malingering, is a stand-alone measure with a 5% false negative rate (Rees, Tombaugh, Gansler, & Moczynski, 1998). The Luria-Nebraska Neuropsychological Battery (LNNB) is a comprehensive neuropsychological test the within-test malingering formula of which has a 17% false negative rate (McKinzey et al., 1997). However, neither is an IQ test, and does not have the same place in a battery as the Raven SPM. More importantly, no test, including the TOMM and LNNB, has published malingering measures validated on children or adolescents (as of this writing).

There are many identifiable groups that were not included in this study. For example, there were no neurologically impaired patients, people with tested IQs below 70, or forensic samples. The current participants are all Austrian, although the Raven is widely used with people speaking a wide range of languages. The faking formula should be cross-validated with such groups in future research, and appropriate interpretive caution employed until such research is done.

### Acknowledgement

The authors would like to thank Brigitte Haider for her help in data collection.

### References

- Faust, D. (1996). Neuropsychological (Brain Damage) Assessment. In J. Ziskin (Ed.), *Coping with psychiatric and psychological testimony* (5th ed., Vol. 2, pp. 916-1044). Los Angeles: Law & Psychology Press.
- Faust, D., Hart, K. J., & Guilmette, T. J. (1988). Pediatric malingering: The capacity of children to fake believable deficits on neuropsychological testing. *Journal of Consulting & Clinical Psychology, 56*(4), 578-582.
- Faust, D., Hart, K. J., Guilmette, T. J., & Arkes, H. R. (1988). Neuropsychologists' capacity to detect adolescent malingerers. *Professional Psychology: Research & Practice, 19*(5), 508-515.
- Faust, D., Ziskin, J., & Hiers, J. (1991). *Brain damage claims: Coping with neuropsychological evidence*. Los Angeles: Law & Psychology Press.
- Gudjonsson, G., & Shackleton, H. (1986). The pattern of scores on Raven's Matrices during "faking bad" and "non-faking" performance. *British Journal of Clinical Psychology, 25*, 35-41.
- Heaton, R. K., Smith, H. H., Lehman, R. A., & Vogt, A. T. (1978). Prospects for faking believable deficits on neuropsychological testing. *Journal of Consulting & Clinical Psychology, 46*(5), 892-900.
- McKinzey, R. K., Podd, M. H., Krehbiel, M. A., Mensch, A. J., & Trombka, C. C. (1997). Detection of malingering on the Luria-Nebraska Neuropsychological Battery: An initial and cross-validation. *Archives of Clinical Neuropsychology, 12*(5), 505-512.
- McKinzey, R. K., Podd, M. H., Krehbiel, M. A., & Raven, J. (1999). Detection of malingering on the Raven Progressive Matrices: A cross-validation. *British Journal of Clinical Psychology, 38*(3), 435-439.
- Raven, J. (2000). *Raven Manual: Research supplement 3* (3rd ed.). Oxford: Oxford Psychologists Press.
- Raven, J., Raven, J. C., & Court, J. H. (2000). *Manual for Raven's Progressive Matrices and Vocabulary Scales. Section 3: The standard progressive matrices*. Oxford: Oxford Psychologists Press.
- Raven, J. C. (1958). *Standard progressive matrices, 1956 revision*. Oxford/San Antonio, TX: Oxford Psychologists' Press/The Psychological Corp., respectively.

- Rees, L. M., Tombaugh, T. N., Gansler, D. A., & Moczynski, N. P. (1998). Five validation experiments of the Test of Memory Malingering (TOMM). *Psychological Assessment, 10*(1), 10-20.
- Rogers, R., & Cruise, K. R. (1998). Assessment of malingering with simulation designs: Threats to external validity. *Law and Human Behavior, 22*(3), 273-285.
- Rogers, R., Harrell, E. H., & Liff, C. D. (1993). Feigning neuropsychological impairment: A critical review of methodological and clinical considerations. *Clinical Psychology Review, 13*(3), 255-274.

Received 1 March 2001; revised version received 21 January 2002